

VIROPHOR 2.8%

DEFRA TB APPROVED STABILISED IODOPHOR DISINFECTANT FOR SURFACES AND FOOT DIP APPLICATIONS

RECOMMENDED USAGE

Remove gross soiling. Apply with a pressure washer or via foam lance at a rate of 300ml/m². Ensure all surfaces are thoroughly wetted.

- For routine surface disinfection use at a dilution rate of 2% v/v of product in dissolved water.
- For foot-dip applications, use at 2% v/v and replace every 2-3 days, or daily if soiling occurs.



For disinfection under the UK control legislation for specific diseases please follow the approved dilution rates shown below;



DEFRA Approvals

Foot and Mouth Disease:	1:550
Swine vesicular Disease:	1:150
Diseases of Poultry (Newcastle):	1:99
Tuberculosis:	1:20
General Orders (Salmonella):	1:50

Storage: Store in tightly closed container in a cool place below 25°C and away from direct sunlight. Store away from foodstuffs and animal feedstuffs.

Pack Size: 4x5, 25 & 200 litre

Shelf Life: 2 years from date of manufacture

USE BIOCIDES SAFELY. FOR HEALTH & SAFETY INFORMATION, REFER TO THE SAFETY DATA SHEET

KILCO (INTERNATIONAL) LTD.
BROOMHOUSES 2 INDUSTRIAL ESTATE
OLD GLASGOW ROAD, LOCKERBIE.
DG11 2SD

Tel: +44 (0) 1576 205 480
Fax: +44 (0) 1576 205 483
Email: enquiries@kilco.co.uk

